

Assessment of Depression, Anxiety and Stress among Patients with Type-II Diabetes Mellitus Study in a Bangladeshi Tertiary Care Hospital

Dr. Mrinal Kanti Sana¹, Dr. Manzurul Ibrahim Musa², Dr. Mahmud Hossain Ibrahim³,
Dr. Hadiur Rahman Siam⁴, Dr. Imtiaz Hasan⁵, Dr. Upananda Roy⁶

¹Assistant Registrar, Department of Respiratory Medicine, Khulna Medical College Hospital, Khulna, Bangladesh.

²Medical Officer, Department of Respiratory Medicine, Khulna Medical College Hospital, Khulna, Bangladesh.

³Registrar, Department of Respiratory Medicine, Khulna Medical College Hospital, Khulna, Bangladesh.

⁴Assistant Registrar, Department of Respiratory Medicine, Khulna Medical College Hospital, Khulna, Bangladesh.

⁵Registrar, Department of Medicine, Khulna Medical College Hospital, Khulna, Bangladesh.

⁶OSD, DGHS, Mohakhali, Dhaka, Bangladesh.

***Corresponding Author:** Dr. Mrinal Kanti Sana, Assistant Registrar, Department of Respiratory Medicine, Khulna Medical College Hospital, Khulna, Bangladesh.

Abstract

Background: Diabetes is a chronic disease caused by insufficient insulin production or ineffective insulin use. The global diabetes population is rapidly increasing, with over 70% living in developing countries. Anxiety and depression often coexist with diabetes, impacting treatment adherence and quality of life. People with diabetes are twice as likely to experience these mental health issues. Understanding and managing the emotional burden of diabetes is crucial for improving overall well-being and treatment outcomes.

Aim of the Study: This study aims to assess the mental health (Depression, anxiety and stress) of patients with Type II Diabetes Mellitus.

Methods: The study conducted at Khulna Medical College and Hospital lasted from 2022 to 2023, with 120 patients enrolled, all with a minimum 2-year history of type 2 diabetes mellitus. Inclusion criteria included patients over 20 years old and male and female patients with a specified diabetic history. Exclusion criteria involved patients with critical health conditions or communication difficulties. Data collection utilized structured and semi-structured questionnaires, as well as chart examinations. Statistical analysis was conducted using SPSS version 26.

Result: The study examined 120 patients, finding that those over 60 years old accounted for 48.33%, with 47.50% aged 40-60 and only 4.17% under 40. Most patients were male (59.17%) and urban residents (79.10%), with 74.17% literate. Mental health issues were prevalent, with 34.17% experiencing stress, 32.50% anxiety, and 33.33% depression. Severity varied: mild group (50% depression, 11.11% anxiety, 38.89% stress), moderate group (56.52% depression, 26.09% anxiety, 17.39% stress), severe group (5.56% depression, 38.89% anxiety, 55.56% stress), and very severe group (15% depression, 55% anxiety, 30% stress).

Conclusion: The study found high rates of depression, anxiety, and stress in Type II Diabetes Mellitus (T2DM) patients, with factors like age, gender, education, and location influencing mental health. Integrating psychological assessments and interventions into T2DM management, routine screening, and multidisciplinary approaches can improve treatment adherence and quality of life.

Keywords: Assessment, depression, anxiety, stress and Type-II DM.

1. INTRODUCTION

Diabetes is a chronic disease that occurs either when the pancreas does not produce enough insulin or when the body cannot effectively use

the insulin it produces [1]. Insulin is a hormone that regulates blood glucose [2]. There are four types of diabetes: Type 1 diabetes is an autoimmune disease. The immune system

attacks and destroys cells in the pancreas, where insulin is made. It is unclear what caused this attack. Type 2 diabetes occurs when your body becomes resistant to insulin, and sugar builds up in your blood. It is the most common type, 90% to 95% Trusted Source of people living with diabetes have type 2 [3], Autoimmune diabetes in adults (LADA). It occurs during adulthood and sets in gradually, like type 2 diabetes. LADA is an autoimmune disease that cannot be treated by diet or lifestyle [4]. Gestational diabetes is high blood sugar during pregnancy [5]. Insulin-blocking hormones produced by the placenta cause this type of diabetes [6]. The number of people affected by diabetes worldwide is increasing rapidly. Currently, it is estimated that nearly 285 million individuals suffer from this condition, and it is projected that this number will increase to 438 million by 2030. More than 70% of people affected by diabetes live in developing countries. [7]. Anxiety and depression affect populations worldwide, but over two-thirds of those affected live in developing countries [8]. The coexistence of Type II Diabetes Mellitus (T2DM) and mental health disorders, specifically depression, anxiety, and stress, constitutes a multifaceted challenge that significantly impacts the well-being of affected individuals [9]. The demands of self-management, dietary restrictions, and potential complications associated with diabetes can lead to heightened levels of emotional distress due to its chronic nature [10]. When not adequately addressed, this emotional burden can adversely affect treatment adherence, glycemic control, and overall quality of life [11]. People with diabetes are almost twice as likely to suffer from anxiety and depression as the general population [12,13]. Recognizing the importance of understanding the psychological dimensions of diabetes, there has been a growing interest in assessing and addressing depression, anxiety, and stress in individuals with Type II Diabetes Mellitus [14]. Despite the evident significance of this interconnection, there remains a need for comprehensive studies that systematically assess the prevalence and impact of depression, anxiety, and stress among individuals with Type II Diabetes Mellitus [15]. This study aims to assess the mental health (Depression, anxiety and stress) of patients with Type II Diabetes Mellitus.

2. METHODOLOGY & MATERIALS

This cross-sectional study was conducted at the Department of Medicine in Khulna Medical

College and Hospital. The study duration was one year, from 2022 to 2023. During the study period, 120 patients were enrolled and analyzed. All patients have at least a 2-year history of type 2 diabetes mellitus.

Inclusion Criteria

- Patients aged more than 20 years.
- Both male and female.
- Patients with at least a 2-year history of type 2 diabetes mellitus.

Exclusion Criteria

- Patients who had critical health conditions (uncertain prognosis, vital signs are unstable, there are major complications).
- Patients were unable to communicate properly.

Data collection involved conducting face-to-face interviews utilizing structured and semi-structured questionnaires and a thorough examination of patients' charts. Sociodemographic information of the participants was gathered through structured questionnaires, which were modified from various sources in the literature. All variables were categorical, with the exception of age, which was obtained through a closed-ended questionnaire format. Statistical analysis Analysis of data was performed using SPSS version 20.0. Categorical variables were expressed as percentages. Comparison between variables was done by using the chi-square test. Association between variables was considered statistically significant if the p-value was less than 0.05.

3. RESULT

This cross-sectional study investigated a cohort of 120 patients, examining various demographic and health factors. The age distribution revealed that 48.33% of the patients were over 60, while 47.50% fell between 40 and 60 years, with only 4.17% under 40. The mean age was 60.25 ± 10.85 years (Table 1). Gender distribution showed a majority of males (59.17%) compared to females (40.83%) (Figure 1). According to Table 2, urban residents comprised 79.10% of the sample, with rural residents comprising 20.83%. Regarding education in Table 3, 74.17% of patients were literate, while 25.83% were illiterate. Analysis of mental health indicators revealed that 34.17% suffered from stress, 32.50% from anxiety, and

33.33% from depression (Table 4). Table 5 presents data on the prevalence of depression, anxiety, and stress across different severity levels among individuals. The severity levels are categorized as mild, moderate, severe, and very severe. In the mild group, comprising 18 individuals, 50.00% reported depression, 11.11% reported anxiety, and 38.89% reported stress. In the moderate group (N=46), 56.52%

reported depression, 26.09% reported anxiety, and 17.39% reported stress. Among those classified as severe (N=36), only 5.56% reported depression, 38.89% reported anxiety, and 55.56% reported stress. Lastly, in the very severe group (N=20), 15.00% reported depression, 55.00% reported anxiety, and 30.00% reported stress.

Table1. Age distribution of the study population (N=120).

Age groups in years	Frequency (n)	Percentage (%)
20-40	5	4.17
40-60	57	47.50
>60	58	48.33
Mean±SD	60.25± 10.85	




Figure1. Gender distribution of the study population (N=120).

Table2. Resident area of the study population (N=120).

Resident	Frequency (n)	Percentage (%)
Urban	95	79.17
Rural	25	20.83
Total	120	100.00

Table3. Education status of the study population (N=120).

Education	Frequency (n)	Percentage (%)
Illiterate	31	25.83
Literate	89	74.17
Total	120	100.00

Table4. Prevalence of depression, anxiety and stress of the study population (N=120).

Variables	Frequency (n)	Percentage (%)
Depression	40	33.33
Anxiety	39	32.50
Stress	41	34.17
Total	120	100.00

Table5. Comparing depression, anxiety and stress with grading.

Variables	Mild (N=18)		Moderate (N=46)		Severe (N=36)		Very severe (N=20)		Total	
	n	%	n	%	n	%	n	%	n	%
Depression	9	50.00	26	56.52	2	5.56	3	15.00	40	33.33
Anxiety	2	11.11	12	26.09	14	38.89	11	55.00	39	32.50
Stress	7	38.89	8	17.39	20	55.56	6	30.00	41	34.17

4. DISCUSSION

This study assessed the depression, anxiety and stress among 120 patients with type 2 diabetes mellitus admitted to a hospital in Khulna, Bangladesh. Depression, anxiety and stress are high among the admitted patients suffering from type 2 diabetes mellitus. Many factors, such as age, current living status, and educational status, are associated with anxiety and depression. Our study scrutinized the age distribution of individuals with Type II Diabetes Mellitus, highlighting that 47.50% of participants were in the 40-60 age range, with almost all patients (48.33%) aged over 60, while the least patients of 4.17% belonged to the 20-40 age group. A parallel study on Type II Diabetes Mellitus by Sharma et al. exhibited similar age patterns, with 47.3% falling in the 40-60 age range and 48.6% aged 60 and above [16]. Nigussil's study demonstrated a broader age range distribution, with the majority falling between 40 and 69 years [17]. These consistent findings across studies reinforce the notion that individuals aged 40 and above are more susceptible to Type II Diabetes Mellitus. In our study, the gender distribution indicates that 59.17% of participants identify as male, while 40.83% identify as female. This gender distribution is consistent with Abualhamael et al. 's study, which demonstrated a more pronounced gender disparity, with 65.7% males and 34.3% females [18]. In contrast to our study, Sharma et al. reported a slightly higher percentage of females (59.5%) compared to females (40.5%) [16]. The unity in gender distribution across these studies underscores a consistent pattern, suggesting that gender ratios in the context of mental health and Type II Diabetes Mellitus are relatively comparable across different populations. In our study, the residential distribution among participants reveals a substantial urban predominance, with 79.17% residing in urban areas and only 20.83% in rural settings. This contrasts with the findings from Sharma et al.'s study, where the urban-resident proportion was 68.6%, and rural residents constituted 31.4% [16]. This urban-centric demographic aligns with the findings from our investigation, emphasizing a concentration of individuals with Type II Diabetes Mellitus in urban settings. Cross-referencing these results with similar investigations could offer a broader understanding of regional variations and their potential impact on mental health outcomes in

individuals with Type II Diabetes Mellitus. Abualhamael et al.'s study demonstrated a diverse educational distribution, with varying levels of formal education among participants, including secondary education (17.9%) and higher education qualifications such as bachelor's degrees (38.2%), master's degrees (8.0%), Ph.D. (6.4%), and diplomas (17.1%) [18]. In contrast, our study presents a simpler classification, with 74.17% of participants considered literate and 25.83% classified as illiterate. In our study, the prevalence of depression, anxiety and stress was 33.33%, 32.5% and 34.17%, respectively, which is comparable with 11.5%, 30.5% and 12.5% among type II diabetic outpatients in Klang Valley, Malaysia [19]. In a similar study done in Pulau Penang and Melange of Malaysia, the prevalence was 26.6%, 40% and 19.4%, respectively [20]. For depression, our findings indicate that 50.00% of cases fall into the mild category, 56.52% are classified as moderate, 5.56% as severe, and 15.00% as very severe, contributing to an overall depression prevalence of 33.33%. A study in Saudi Arabia reported rates of mild, moderate, severe, and extremely severe depression among patients with type 2 DM as 9.3%, 14.0%, 7.1%, and 3.3%, respectively [21]. Another study found that 12.4% of participants experienced mild depression, 29.7% had moderate depression, 3.4% had severe depression, and 4.2% had very severe depression, resulting in an overall depression prevalence of 49.7% [22]. In terms of anxiety, 11.11% are categorized as mild, 26.09% as moderate, 38.89% as severe, and 55.00% as very severe, resulting in an overall anxiety prevalence of 32.50% according to our study. In the same line, a study in Saudi Arabia showed mild, moderate, severe and extremely severe anxiety among 13.4, 13.0%, 6.0%, and 5.8% of patients with type 2 DM, respectively [21]. However, a very low prevalence of anxiety was observed in other studies conducted in 15 nations and in the United States, Baltimore, which showed 18.0% and 21.8% overall prevalence of anxiety, respectively, among type 2 diabetes patients [23,24].

Lastly, the distribution for stress shows 38.89% in the mild category, 17.39% in the moderate range, 55.56% as severe, and 30.00% as very severe, contributing to an overall stress prevalence of 34.17% in our study. A similar study reported that their distribution indicates

11.0% with mild stress, 18.6% with moderate stress, 27.6% with severe stress, and 11.0% with severe stress, resulting in an overall stress prevalence of 68.3% [22]. The variation in the results might be due to the nature of the patients included in these studies and their different sociodemographic characteristics and measurement tools.

Limitations of the Study: The study's sample size of 120 patients from a single Khulna, Bangladesh hospital may not represent the broader population of individuals with Type II Diabetes Mellitus. The findings may not generalise to other settings or populations with different demographic characteristics and healthcare systems. Moreover, the exclusion of patients with critical health conditions may have biased the results, as these individuals may experience higher levels of depression, anxiety, and stress. Finally, the assessment of depression, anxiety, and stress relied on self-report measures, which may be subject to social desirability bias and may not accurately reflect patients' mental health status.

5. CONCLUSION AND RECOMMENDATIONS

In conclusion, our study highlights a significant prevalence of depression, anxiety, and stress among patients with Type II Diabetes Mellitus (T2DM). Factors such as age, gender, educational status, and residential area show associations with mental health indicators. The high prevalence of these mental health disorders underscores the importance of integrating psychological assessments and interventions into the management of T2DM. Recommendations include routine screening for depression, anxiety, and stress in T2DM patients during clinical visits, as well as the implementation of multidisciplinary approaches involving healthcare providers, psychologists, and social workers to address the psychological well-being of individuals with T2DM. Early identification and management of mental health issues can improve treatment adherence, glycemic control, and overall quality of life for T2DM patients.

REFERENCES

[1] Mukhtar Y, Galalain A, Yunusa U. A modern overview on diabetes mellitus: a chronic endocrine disorder. *European Journal of Biology*. 2020 Nov 23; 5(2):1-4.

[2] Qaid MM, Abdelrahman MM. Role of insulin and other related hormones in energy metabolism—A review. *Cogent Food & Agriculture*. 2016 Dec 31; 2(1):1267691.

[3] Trikkalinou A, Papazafiroπούλου AK, Melidonis A. Type 2 diabetes and quality of life. *World journal of diabetes*. 2017 Apr 4; 8(4):120.

[4] Buzzetti R, Tuomi T, Mauricio D, Pietropaolo M, Zhou Z, Pozzilli P, Leslie RD. Management of latent autoimmune diabetes in adults: a consensus statement from an international expert panel. *Diabetes*. 2020 Oct 1; 69(10):2037-47.

[5] Palani S, Joseph NM, Tegene Y, Zacharia A, Marew T. Gestational diabetes-A review. *JGTPS*. 2014;5(2):1673-83.

[6] Mukhtar Y, Galalain A, Yunusa U. A modern overview on diabetes mellitus: a chronic endocrine disorder. *European Journal of Biology*. 2020 Nov 23;5(2):1-4.

[7] Soedamah-Muthu SS, Abbring S, Toeller M. Diet, lifestyle and chronic complications in type 1 diabetic patients. In *Type 1 Diabetes-Complications, Pathogenesis, and Alternative Treatments 2011* (pp. 21-44). InTech.

[8] Khuwaja AK, Qureshi R, Azam SI. Prevalence and factors associated with anxiety and depression among family practitioners in Karachi, Pakistan. *Journal of Pakistan medical association*. 2004; 54(2):45.

[9] Randväli M, Toomsoo T, Šteinmiller J. The Main Risk Factors in Type 2 Diabetes for Cognitive Dysfunction, Depression, and Psychosocial Problems: A Systematic Review. *Diabetology*. 2024 Jan 11;5(1):40-59.

[10] Kalra S, Jena BN, Yeravdekar R. Emotional and psychological needs of people with diabetes. *Indian journal of endocrinology and metabolism*. 2018 Sep; 22(5):696.

[11] Weinger K, Jacobson AM. Psychosocial and quality of life correlates of glycemic control during intensive treatment of type 1 diabetes. *Patient education and counseling*. 2001 Feb 1; 42(2):123-31.

[12] Nichols L, Barton PL, Glazner J, McCollum M. Diabetes, minor depression and health care utilization and expenditures: a retrospective database study. *Cost Effectiveness and Resource Allocation*. 2007 Dec;5(1):1-8.

[13] Bouwman V, Adriaanse MC, van't Riet E, Snoek FJ, Dekker JM, Nijpels G. Depression, anxiety and glucose metabolism in the general dutch population: the new Hoorn study. *PLoS one*. 2010 Apr 1; 5(4):e9971.

[14] Kalra S, Jena BN, Yeravdekar R. Emotional and psychological needs of people with diabetes. *Indian journal of endocrinology and metabolism*. 2018 Sep; 22(5):696.

[15] Basiri R, Seidu B, Rudich M. Exploring the Interrelationships between Diabetes, Nutrition, Anxiety, and Depression: Implications for

- Treatment and Prevention Strategies. *Nutrients*. 2023 Sep 30; 15(19):4226.
- [16] Sharma K, Dhungana G, Adhikari S, Bista Pandey A, Sharma M. Depression and anxiety among patients with type ii diabetes mellitus in Chitwan Medical College Teaching Hospital, Nepal. *Nursing research and practice*. 2021 Jan 13; 2021.
- [17] Nigussie K, Sertsu A, Ayana GM, Dessie Y, Bete T, Abdisa L, Debele GR, Wadaje D, Negash A. Determinants of depression and anxiety among type 2 diabetes patients in governments' hospitals at Harari regional state, Eastern Ethiopia: A multi-center cross-sectional study. *BMC psychiatry*. 2023 Jan 5;23(1):13.
- [18] Abualhamael SA, Baig M, Alghamdi W, Gazzaz ZJ, Al-Hayani M. Quality of life, stress, anxiety and depression and associated factors among people with type 2 diabetes mellitus in Western region Saudi Arabia. *Frontiers in Psychiatry*. 2024 Jan 15; 14:1282249.
- [19] Kaur G, Tee GH, Ariaratnam S, Krishnapillai AS, China K. Depression, anxiety and stress symptoms among diabetics in Malaysia: a cross sectional study in an urban primary care setting. *BMC family practice*. 2013 Dec; 14:1-3.
- [20] Tan KC, Chan GC, Eric H, Maria AI, Norliza MJ, Oun BH, Sheerine MT, Wong SJ, Liew SM. Depression, anxiety and stress among patients with diabetes in primary care: a cross-sectional study. *Malaysian family physician*: the official journal of the Academy of Family Physicians of Malaysia. 2015; 10(2):9.
- [21] Alzahrani A, Alghamdi A, Alqarni T, Alshareef R, Alzahrani A. Prevalence and predictors of depression, anxiety, and stress symptoms among patients with type II diabetes attending primary healthcare centers in the western region of Saudi Arabia: a cross-sectional study. *International journal of mental health systems*. 2019 Dec;13(1):1-7.
- [22] Niraula K, Kohrt BA, Flora MS, Thapa N, Mumu SJ, Pathak R, Stray-Pedersen B, Ghimire P, Regmi B, MacFarlane EK, Shrestha R. Prevalence of depression and associated risk factors among persons with type-2 diabetes mellitus without a prior psychiatric history: a cross-sectional study in clinical settings in urban Nepal. *BMC psychiatry*. 2013 Dec; 13(1):1-2.
- [23] Chaturvedi SK, Gowda SM, Ahmed HU, Alosaimi FD, Andreone N, Bobrov A, Bulgari V, Carrà G, Castelnuovo G, de Girolamo G, Gondek T. More anxious than depressed: prevalence and correlates in a 15-nation study of anxiety disorders in people with type 2 diabetes mellitus. *General Psychiatry*. 2019; 32(4).
- [24] Edwards LE, Mezuk B. Anxiety and risk of type 2 diabetes: evidence from the Baltimore Epidemiologic Catchment Area Study. *Journal of psychosomatic research*. 2012 Dec 1; 73(6):418-23.

Citation: Dr. Mrinal Kanti Sana et al. *Assessment of Depression, Anxiety and Stress among Patients with Type-II Diabetes Mellitus Study in a Bangladeshi Tertiary Care Hospital*. *ARC Journal of Diabetes and Endocrinology*. 2024; 9(1):1-6. DOI: <http://dx.doi.org/10.20431/2455-5983.0901001>.

Copyright: © 2024 Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.