

The Mystery of the Mirror Neuron System

Eden Wudneh¹, Ajita Acharya², Aisha Ashraf³, Radhika Krishnan⁴, Hassaan Tohid^{5,6}

^{1,2,3,4,5}Department of Neurology, California Institute of Behavioral Neurosciences & psychology
(The Neuro-Cal Institute)

⁶Center for Mind & Brain, University of California, Davis

Abstract:

Objective: *To highlight the link of mirror neurons with body movements*

Method: *A literature search was conducted in PubMed and various other data bases to collect some studies about the association of mirror neurons with body movement.*

Result: *Studies do exist that support the notion that there are some sets of neurons in the human and animal brain that are activated when humans or animals perform certain actions. The same neurons are activated when they observe another person or animal performing some movement like walking and hand related movement.*

Conclusion: *The area of the human and animal brain that is activated when we perform an action and when we see someone else performing the same action, is believed to have a special kind of neurons known as “mirror neurons”. The evidence proving the existing of these new neurons is still not sufficient, as these neurons were discovered just couple of decades ago. We believe future studies will be useful to provide more information about the subject.*

Keywords: *Mirror Neurons, Mirror Neurons legs, mirror neurons limbs, mirror neurons hands, mirror neurons grip, mirror neurons movement, mirror neurons motion.*

1. BACKGROUND

According to the world renowned mirror neurons expert Dr. VS Ramachandran, “I predict that mirror neurons will do for psychology what DNA did for biology” (1). Sounds strange, but if we study mirror neurons in depth we will find out that this new discovery of mirror neurons has a potential to unearth so many mysteries, like why we yawn, when we see the other person yawning, and why we smile when we see other person smiling, or why we feel sad when seeing a sad person (2).

Mirror neurons are a distinct group of neurons that fire both when an individual executes a motor act and when he/she observes another individual performing the same or a similar motor act (3-4). Thus the neuron “mirrors” the behavior of the other. Mirror neurons were first discovered over 20 years ago in monkey’s brain, in the inferior frontal gyrus (area F5) and inferior parietal gyrus. As of now, these neurons have been found in the premotor cortex, the supplementary motor area, the inferior parietal cortex, and the primary somatosensory cortex (5).

Many new readers who read about mirror neurons have a tendency to believe that mirror neurons are only related to facial expressions or emotions. However, recent research and observations have broadened our understanding about these unique neurons. Today we do know that these neurons are associated with many bodily functions.

Not much was known about the association of mirror neurons with body movements and function until the extensive fMRI studies made it possible to dig deeper into the human and animal brain and extract the answers to this unsolved mystery. In this article, we are not hypothesizing any question, but we are only bringing together some recent pieces of evidence regarding the association of mirror neurons with various other body functions especially movement.

2. DISCUSSION

During our various activities of daily living, we communicate, socialize and connect with other people and we somehow recognize the state of mind they are in and what they will do or how they will likely act. Different motor simulation theories of perception state that effective perceptual processing of others' actions can be attained not solely by visual analysis of the movements but also by a process of

motor simulation. To disprove this notion, Vannuscorps et al conducted eight experiments and found that individuals born with absent or severely shortened upper limbs (upper limb dysplasia), could understand, anticipate and recall upper limb movements, which they cannot mimic, as efficiently as typically developed participants [6].

The function of mirror neurons has been, in some way, elucidated by the method coaches use to train athletes. They apply a technique called “action demonstration” to make athletes acquire new motor tasks [7,8]. To achieve this, the learners are presented with a model that demonstrates how to perform the action they want to learn [9, 10]. The learners then need to transform the observed visual information into motor commands that allow them to execute the action correctly (visio-motor transformation)[11,12]. In these trainings two apprentices work together. Alternating the physical practice with one another, the second player observes and executes what the first player does. Then the first player observes the second player and executes the same thing [13]. The observation of his/her partner and the subsequent execution of the task would be the factors that increase performance efficiency.

A study by Salvia et al proposed that observing others' actions augments muscle-specific cortico-spinal excitability, indicating presumed mirror neurons activity. They also suggested that the exposure to emotional stimuli modify cortico-spinal excitability. The interaction of these two phenomena when they are combined was further explored. A single-pulse transcranial magnetic stimulation (TMS) was delivered over the hand area of the left primary motor cortex of 27 healthy adults and adolescents and their right first dorsal interosus (FDI) muscle activity (i.e., motor evoked potential - MEP) was recorded, while they watched either videos of neutral or angry hand actions and facial expressions, or neutral objects as a control condition. They were able to recreate the motor resonance and the emotion effects. It was found that hand-actions and emotional stimuli triggered greater cortico-spinal excitability than the faces/control condition and neutral videos, respectively. Furthermore, the impact of emotion was present for faces but not for hand actions, which indicated that the motor resonance and the emotion effects might not be additive [14].

Furthermore, Maeda et al emphasized the important role of parieto-frontal pathways in visually guided motor control. They documented 235 neurons related to hand manipulation exercises. Of these, fifty-four showed the same response to both the video clips of the monkey's own hand movement as well as to the visual feedback during that action or clips of the experimenter's hand movement in a lateral view. Twenty-five out of the 54 were labeled “hand-type” as they only responded to video clips of the monkey's own hand, without an image of the target object. Thirty-three of 54 neurons that were defined as mirror neurons showed visual responses to the experimenter's action and motor responses. The study concluded that activity of hand manipulation-related and mirror neurons in anterior intraparietal/Parieto-frontal gyrus (PFG) plays a fundamental role in monitoring one's own body state based on visual feedback [15].

The effect of mirror neurons on empathy has been shown in different studies. The same neural circuits that fire when a person is happy are also activated when we see someone is feeling happy [16]. Another way of exploring the role of mirror neurons in empathy is through pain and disgust [17]. It was found that applying painful stimuli to subject's hands activated neurons in the anterior cingulate cortex. Similarly, the same neurons discharged when watching painful stimuli being applied to someone else's hands. The neurons in the anterior cingulate gyrus also fired when noticing people who would be painfully stimulated at a later time. Hence, the area of the brain accounted for reacting to pain are activated not only while experiencing pain but also watching someone else experiencing pain as well as noticing someone else who would encounter pain at a later time [18]. Similarly, neurons in the Insula, which fire when the person experiences disgust, also become activated when he/she observes faces showing disgust. This shows that when a person is incapable of experiencing emotion, this person will likely have impaired understanding of the same emotions in others. A case report written on a patient with brain injury entailed that damage to the area of the brain responsible for experiencing disgust led to difficulty detecting disgust in others [17].

Another concept coined about mirror neurons' function is Mirror-touch synesthesia. This is a condition where seeing other individuals experiencing somatosensory sensation causes the same sensation in the observer [19]. The most supported theory behind this phenomenon is that in normal people the somatosensory mirror system, which moderate observed and felt touch, has activations that are below a certain threshold. When the activations are below this threshold, a person can sense and

appreciate the observed touch. In motor neuron synesthesia the mental imitation is so strong that it crosses a threshold into near-tactile sensation, which is sometimes indistinguishable from the synesthete's own [20]. This condition has been seen in patients who have suffered stroke with loss of sensation and amputees where visual stimuli elicited a tactile sensation [21].

3. CONCLUSION

The mirror neuron system (MNS) is linked with many functions including the body movements such as limb movements such as walking, hand movement like gripping and grasping etc. Action observation is also found associated with mirror neurons. It is also found that observing others' actions augments muscle-specific cortico-spinal excitability, indicating presumed mirror neurons activity. Empathy and mirror-touch synesthesia has also been studied in relation to mirror neurons.

There so much more to mirror neurons. This article is just a mini review to highlight some facts about the mirror neurons, to help new scientists and clinicians learn some basics about mirror neuron system. Although it has been more than two decades since the mirror neurons were first discovered, the data present is still not sufficient and future research studies will help us learn more about these mysterious kinds of neurons.

REFERENCE

- [1] Ramachandran V. MIRROR NEURONS and imitation learning as the driving force behind the great leap forward in human evolution. *Conversation: Mind. Edge*. May, 31st, 2000. Available from <https://www.edge.org/conversation/mirror-neurons-and-imitation-learning-as-the-driving-force-behind-the-great-leap-forward-in-human-evolution>. Accessed Aug 24th, 2016.
- [2] Rizzolatti, Giacomo; Craighero, Laila (2004). "The mirror-neuron system" (PDF). *Annual Review of Neuroscience*. 27 (1): 169-192. doi:10.1146/annurev.neuro.27.070203. 144230. PMID 15217330
- [3] Keysers, Christian (2010). "Mirror Neurons" (PDF). *Current Biology*. 19 (21): R971–973. doi:10.1016/j.cub.2009.08.026. PMID 19922849
- [4] Rizzolatti, Giacomo; Fadiga, Luciano (1999). "Resonance Behaviors and Mirror Neurons". *Italianes de Biologie*. 137: 85–100.
- [5] Molenberghs P, Cunnington R, Mattingley J (July 2009). "Is the mirror neuron system involved in imitation? A short review and meta-analysis.". *Neuroscience & Biobehavioral Reviews*. 33 (1): 975–980. doi:10.1016/j.neubiorev.2009.03.010
- [6] Vannuscorps G1, Caramazza A2. Typical action perception and interpretation without motor simulation. *Proc Natl Acad Sci U S A*. 2016 Jan 5; 113(1):86-91. doi: 10.1073/pnas.1516978112. Epub 2015 Dec 22.
- [7] Bandura, A. (1986). *Social foundations of thought and action: a social cognitive theory*. Englewood Cliffs, N.J.: Prentice-Hall.
- [8] McCullagh, P., & Weiss, M. R. (2001). Modeling: considerations for motor skill performance and psychological responses. *Handbook of sport psychology*, 205-238.
- [9] Al-Abood, S. A., Davids, K., Bennett, S. J., Ashford, D., & Martinez Marin, M. Effects of manipulating relative and absolute motion information during observational learning of an aiming task. (2001). *J Sports Sci*, 19(7), 507-520.
- [10] Buchanan, J. J., & Dean, N. J. (2010). Specificity in practice benefits learning in novice models and variability in demonstration benefits observational practice. *Psychol Res*, 74(3), 313-326.
- [11] Jeannerod, M., Arbib, M. A., Rizzolatti, G., & Sakata, H. (1995). Grasping objects: the cortical mechanisms of visuomotor transformation. *Trends Neurosci*, 18(7), 314-320.
- [12] Rizzolatti, G., Luppino, G., & Matelli, M. (1998). The organization of the cortical motor system: new concepts. *Electroencephalogr Clin Neurophysiol*, 106(4), 283-296.
- [13] Wulf G, Shea C, Lewthwaite R. Motor skill learning and performance: a review of influential factors. *Med Educ*. 2010 Jan; 44(1):75-84. doi: 10.1111/j.1365-2923.2009.03421.x.
- [14] Salvia E1, Süß M2, Tivadar R2, Harkness S2, Grosbras MH3. Mirror Neurons System Engagement in Late Adolescents and Adults While Viewing Emotional Gestures. *Front Psychol*. 2016 Jul 20; 7:1099. doi: 10.3389/fpsyg.2016.01099. eCollection 2016.

- [15] Maeda K1, Ishida H, Nakajima K, Inase M, Murata A. Functional properties of parietal hand manipulation-related neurons and mirror neurons responding to vision of own hand action. *J Cogn Neurosci*. 2015 Mar; 27(3):560-72. doi: 10.1162/jocn_a_00742. Epub 2014 Oct 14
- [16] Banissy, Michael; Jamie Ward (July 2007). "Mirror Touch Synaesthesia is linked with Empathy". *Nature Neuroscience*. 10 (7): 815–816. doi:10.1038/nn1926. PMID 17572672.
- [17] Corradini, Antonella; Alessandro Antonietti (2013). "Mirror neurons and their function in cognitively understood empathy". *Consciousness and Cognition*. 22 (3): 1152–1161. doi:10.1016/j.concog.2013.03.003.
- [18] Singer, T.; B. Seymour; J. O'Doherty; H. Kaube; R.J. Dolan; C.D. Frith (2004). "Empathy for pain involves the affective but not the sensory components of pain". *Science*. 1 (303): 1157–1162. Bibcode:2004Sci...303.1157S. doi:10.1126/science.1093535.
- [19] Banissy, M.J. (2009). "Prevalence, Characteristics, and a Neurocognitive Model of Mirror Touch Synesthesia". *Experimental Brain Research*. 192 (2): 261–272. doi:10.1007/s00221-009-1810-9.
- [20] Blakemore, SJ; D. Bristow; G. Bird; C. Frith; J.Ward (2005). "Somatosensory Activations During the Observation of Touch and a Case of Vision-Touch Synaesthesia". *Brain*. 1 (28): 1571–1583. doi:10.1093/brain/awh500. PMID 15817510.
- [21] Goller, A.I. (2013). "Mirror Touch Synesthesia in the Phantom Limbs of Amputees". *Cortex*. 49 (1): 243–251. doi:10.1016/j.cortex.2011.05.002.