

Different Nacl Concentrations Impact on Seed Germination of Basil (Ocimum Basilicum L.) and Fennel (Foeniculum VulgareL.) Cultivars

Cenk PASA*

Balıkesir University, Altınoluk Vocational School, Deparments of Plant and Animal Production, Medicinal and Aromatical Plant Programme, Altınoluk, Edremit, Balıkesir.

*Corresponding Author: Cenk PASA, Balikesir University, Altinoluk Vocational School, Deparments of Plant and Animal Production, Medicinal and Aromatical Plant Programme, Altinoluk, Edremit, Balikesir.

Abstract

In this research, has been studied the effects of NaCl of different concentrations on all characteristics of germination of basil (Ocimum basilicum L.) and fennel (Foeniculum vulgare). Basil and Fennel varieties have been studied under four salinity treatments (0, 50, 100 and 150 mM) in the laboratory conditions. In the experiments, plant fresh weight, germination rate, hypocotyl and radicula length for all genotypes were decreased at the increase of salt dose. The results show that maximum germination speed of 0 mM (96.70 %) was observed in a salt dose of Basil. The results pointed out that maximum germination speed of 0 mM (94.70 %) was observed in a salt dose at Fennel.

Keywords: Foeniculum vulgare, Ocimum basilicum, NaCl, germination, seed.

1. INTRODUCTION

Throughout history, environmental stress due to the high concentration of salt ions in soils has been one of the most important factors limiting the productivity of agricultural crops, especially plants susceptible to soil salinity. Today, the high salinity level in agricultural soils adversely affects agricultural production in the world1.

The period in which the plants are the most sensitive to salinity is the germination period. Germination on salty soils cannot be sufficient. The reason for this is that salt solutions in the soil move from bottom to top by evaporation2.

Salinity affects the germination rate, percentage and seedling growth in different ways depending on the plant species3,4,5. High levels of salinity affect seed germination and plant growth by water deficit, ion toxity and ion imbalance or a combination of these factors 6,7.

Ocimum (Basil) species are fragrant and perennial plants belonging to Lamiaceae family can grow in a wide area of the world8,9. Basil is used as a spice and has a unique smell due to its essential oils. Many plant species belonging to the Lamiaceae family can grow wild in the Mediterranean region 10,11,12,13,14.

Foeniculum vulgare (Fennel) is a Mediterranean plant used in traditional medicine and as a spice. The fruit of fennel plant can be used as antioxidant, diuretic, analgesic15. Essential oils are mainly concentrated in the pericarps and provide the unique aroma and taste16. It also increases milk flaw and libido and relives menopausal symptoms in women17.

The aim of this research was to characterize the effect of salinity on seed germination of Ocimum basilicum and Foeniculum vulgare cultivars and it has been studied under four salinity treatments, including, 0, 50, 100 and 150 mM in the laboratory conditions. In the experiments, germination rate, plant fresh weight, hypocotyl length and radicula length for all genotypes were decreased at the increase of the salt dose.

2. MATERIAL AND METHODS

O. basilicum and F. vulgare cultivars were provided by the Altinoluk Vocational School, Deparments of Plant and Animal Production, Medicinal and Aromatical Plant Programme, Balikesir University, Turkey. Seeds were sown on sterilized filter paper placed in petri dishes (6 mm). It was then wetted with a solution of different salinity concentrations (0, 50, 100 and 150 mM) as well as distilled water. The study was conducted in four replications. After 4 days germination rate of seeds gives germinating rate; on the other hand, at the end of 8 days, it gives germination power. Germination rate, plant fresh weight, hypocotyl and radicle lengths were evaluated using ten seedlings. Experimental data were analyzed by ANOVA and compared with the Least Significant Difference (LSD) test (p<0.05). For all parameters investigated, variance analysis was performed using MSTAT-C software.

3. RESULTS AND DISCUSSION

The results of this study reveal that salinity concentration characteristics significantly affected all characters (p<0.05). The differences between the means were compared by Duncan's multiple range tests and are shown in Table 1. The results show that maximum germination speed of 0 mM (96.70 %) was observed in a salt dose. The lowest germination speed of 150 mM (88.75 %) was obtained in salt dose. The lowest germination rate of 150 mM (93.50 %); radicula length of 150 mM (0.053 cm %); hypocotyl length rate of 150 mM (0.08 cm %) and plant fresh weight of 150 mM (0.008 g %) were obtained in salt dose. The results show that maximum germination rate of 0 mM (100 %); radicula length of 0 mM (0.96 cm); maximum hypocotyl length of 0 mM (0.24 cm) and plant fresh weight of 0 mM (0.019 g) were observed in a salt dose.

Salt Dose	Germination Speed (%)	Germination Rate (%)	Radicula length (cm)	Hypocotyl length (cm)	Plant fresh weight (g)
0	96.70a*	100.00a	0.96a	0.24a	0.019a
50	95.30b	98.00b	0.88b	0.22ab	0.016b
100	93.10c	94.00c	0.71c	0.15b	0.011c
150	88.75d	93.50c	0.53d	0.08a	0.008d
Mean	93.46	96.38	0.77	0.17	0.014
LSD	1.24	0.97	0.07	0.018	0.002
CV	7.86	6.55	8.81	10.14	9.84

Table 1. Physical measurements of O. basilicum seeds during germination

*Differences among the parameters are statistically significant at p < 0.05

There is no statistically significant difference (p < 0.05) between figures including the same letters in the columns

The results of this study showed all characters were significantly affected by different NaCl concentrations (p<0.05). The differences between these tools were compared with Duncan's multi-class test (Duncan's test) and presented in Table 2.

Salt Dose	Germination Speed (%)	Germination Rate (%)	Radicula length (cm)	Hypocotyl length (cm)	Plant fresh weight (g)
0	94.30a*	100.00a	2.31a	0.53b	0.034a
50	92.80b	97.50b	2.08b	0.64a	0.029b
100	92.10b	96.30c	1.24c	0.31c	0.021c
150	89.30c	93.40d	0.91d	0.14d	0.013d
Mean	92.13	96.80	1.64	0.41	0.024
LSD	1.07	1.13	0.15	0.09	0.0038
CV	6.21	7.02	5.89	7.14	8.86

Table 2. Physical measurements of F. vulgare seeds during germination

* Differences among the parameters are statistically significant at p < 0.05

There is no statistically significant difference (p>0.05) between figures including the same letters in the columns

The results show that maximum germination speed of 0 mM (94.70 %); germination rate of 0 mM (100 %); radicula length of 0 mM (2.31 cm); hypocotyl length of 50 mM (0.64 cm) and plant fresh weight of 0 mM (0.034 g) were observed in a salt dose. The lowest germination speed of 150 mM

International Journal of Advanced Research in Botany (IJARB)

(89.30%);germination rate of 150 mM (93.40%); radicula length of 150 mM (0.91 cm%); hypocotyl length rate of 150 mM (0.14 cm) and plant fresh weight of 150 mM (0.013 g) were obtained in salt dose.

Similar results were found by Maas and Hoffman18 and Huggen19. High salt concentration in germination medium can reduce or delay germination percentage20,21,22. Some plants showed different resistance to salinity during germination23. Under saline conditions, shoot growth was frequently inhibited more than root growth24. Salinity stress had remarkable effects on other plant growth parameters such as plant and root fresh weight25. High foliar concentration of Na+ is capable of reducing CO2 assimilation because of ionic toxicity26. Reduction in plant and root fresh weight in response to salt stress has been reported for other crops, such as okra, broadbean, soybean chickpea, cowpea, black cumin melon, tomato and watermelon27-34.

4. CONCLUSION

The results show that maximum germination speed of 0 mM (96.70 %) was observed in a salt dose of Basil. The results pointed out that maximum germination speed of 0 mM (94.70 %) was observed in a salt dose at Fennel. Results showed that O. basilicum and F. vulgare all characters were depressed when we increased NaCl concentration. Although further investigations are needed to ascertain the present result, some conclusions may be drawn from these findings.

REFERENCES

- [1] H. Zhang and J. Hodson, Engineering salt tolerance Brassica plants: characterization of yield and seed oil quality in transgenic plants with increased vacuolar sodium accumulation, PNAS, 48, 2001, 12832-12836.
- [2] L. Bernstein, Crop growth and salinitiy. p 39-54. In: J. van Scdiffgard Drainage for agriculture. Agroomy. Monography, 17, 1974, ASA, Nadison, WI.
- [3] B. Murillo-Amador, E. Troyo-Dieguez, H.G. Jones, F. Ayala-Chairez, C.L. Tinoco-Ojanguren, A. Lopez-Cortes, Screening and classification of cowpea genotypes for salt tolerance during germination. Phyton Int. J. Exp. Bot., 2000, 67: 71-84.
- [4] M. Almansouri, J.M. Kinet, S. Lutts, Effect of salt and osmotic stresses on germination in durum wheat (Triticum durum Desf.). Plant Soil, 2001, 231: 243-254.5.
- [5] A. El-Keblawy, A. Al-Rawai, Effects of salinity, temperature and light on germination of invasive Prosopis juliflora (Sw.) D.C. J. Arid Environ., 2005, 61: 555-565.
- [6] A. Läuchli, S.R. Grattan, Plant Growth and Development under Salinity Stress Advances, 2007, 1-32 p. In: Jenks MA et al. (Eds.). Molecular Breeding toward Drought and Salt Tolerant Crops. pp. 285-315. (Eds.): M.A. Jenks, P.M. Hasegawa and S.M. Jain. Springer, Dordrecht, Netherlands.
- [7] D.H. Reinhardt, T.L. Rost, On the correlation of the primary root growth and treachery element size and distance from the tip in cotton seedlings grown under salinity. Environmental and Experimental Botany, 1995, 35:575-588.
- [8] A. Akgül, Spice Science and Technology. Turkish Association Food Technologists Publ. No. 15, Ankara, Turkey. (in Turkish), 1993,
- [9] O. Baritaux, H. Richard, J. Touche, M. Derbesy, Effects of drying and storage of herbs and spices on the essential oil. Part I, 1992.
- [10] A. Akgül, Volatile oil composition of sweet basil (Ocimum basilicum L.) cultivating in Turkey. Nahrung, 1989, 33: 87–88.
- [11] M. Marotti, R. Piccaglia, E. Giovanelli, Differences in essential oil composition of basil (Ocimum basilicum L.) Italian cultivars related to morphological characteristics. J. Agr. Food Chem., 1996,14: 3926–3929.
- [12] K. Sanda, K. Koba, P. Nambo, A. Gaset, Chemical investigation of Ocimum species growing in Togo. Flavour Fragr. J.,1998, 13: 226–232.
- [13] A.P. Martins, L.R. Salgueiro, R. Vila, F. Tomi, S. Canigueral, J. Casanova, A.P. Da Cunha, T. Adzet, Composition of the essential oils of Ocimum canum, O. gratissimum and O. minimum. Planta Med., 1999, 65: 187–189.
- [14] Ö. Musa and Jean-Clause-Chalcat, Essential Oil Composition of Ocimum basilicum L. and Ocimum minimum L. in Turkey. Czech J. Food Sci., 2002, Vol. 20, No. 6: 223–22815. E.M. Choi, J.K. Hwang, Antiinflammatory analgesic and antioxidant activities of the fruit of Foeniculum vulgare. Fitoterapia. 2004;75:557–565.

- [15] M. Siahi, A. Shiravi, M. Heydari, Effect of fennel hydro extract of fennel on prolactin and lactation in female Rat. J Damghan, 2009; 1:55–63.
- [16] B. Khani, F. Mehrabian, E. Khalesi, A. Eshraghi, Effect of soy phytoestrogen on metabolic and hormonal disturbance of women with PCOS. J Res Med Sci. 2011;16:297–230.
- [17] E.V. Maas and G.J. Hoffman, Crop salt tolerance-current assessment Journal of Irrigation and Division, 1977, 103, 115-134.
- [18] A. Huggen, Salt tolerance through increased vigor in a flax line (STS-II) elected for salt tolerance in vitro. Theoretical and Applied Genetics, 1987,74, 727-732.
- [19] Y. Waisel, Biology of halophytes. Academic Press, 1972, New York London.
- [20] S. Mohammed and D.N. Seeni, Germination behavior some halophytes in Indian Desert Journal of Experimental Biology, 1990, 28, 545-549.
- [21] M.O. Basalah, Effect of salinity on germination and growth of sequash (Cucurbita pepo L.) Arabian Research, 1991, 25-29.
- [22] J.A. Franco, J.A. Fernandez, S. Banon and A.Gonzalez, Relationship between the effects of salinity on seedling growth and yield of six muskmelon cultivars. HortScience, 1997, 32, 642–644.
- [23] A. Poljakoff-Mayber and H.R. Lerner, Plants in saline environments. In: Handbook of Plant and Crop Stress. (Pessarakli, M., Ed.). Marcel Dekker Inc., New York, NY, USA, 1994, 65–96.
- [24] K. Aydinşakir, B. Büyüktaş, N. Dinç, C. Karaca, Impact of salinity stress on growing, seedling development and water consumption of peanut (Arachis hypogaea cv. NC-7). Akdeniz Üniversitesi Ziraat Fakültesi Dergisi, 2015, 28(2):77-84.
- [25] P. Cachorro, A. Ortiz, A. Cerdá, Effects of saline stress and calcium on lipid composition in bean roots. Phytochemistry, 1993, 32:1131–1136.
- [26] S. De Pascale, G. Barbieri, Effects of salinity and top removal on growth and yield of broadbean as a green vegetable. Scientia Horticultureae, 1997, 71:147-165.
- [27] P.H. Zaidi, B.B. Sing, Dry matter partitioning and yield attributes of soybean as affected by soil salinity and growth regulators. Legume Research, 1993, 16:3-4.
- [28] O. Düzdemir, A. Ünlükara, A. Kurunç, Response of cowpea (Vigna unguiculata) to salinity and irrigation regimes. New Zeland Journal of Crop and Horticultural Science, 2009, 37:271-280.
- [29] A.S. Hajar, M.A. Zidan, H.S. Al-Zahrani, Effect of salinity stress on the germination, growth and some physiological activities of black cumin (Nigella sativa L.). Arab Gulf Journal of Scientific Research, 1996, 14(2): 445-454.
- [30] H.O. Sivritepe, N. Sivritepe, A. Eris E. Turhan, The effects of NaCl pre-treatments on salt tolerance of melons grown under long-term salinity. Scientia Horticulturae, 2005, 106(4):568-581.
- [31] A. Ünlükara, A. Kurunç, D.G. Kesmez, E. Yurtseven, Effects of salinity on eggplant (Solanum melongena L.) growth and evapotranspiration. Irrigation and Drainage, 2008, 59:203-214.
- [32] E. Yurtseven, G.D. Kesmez, A. Ünlükara, The effects of water salinity and potassium levels on yield, fruit quality and water consumption of a native central Anatolian tomato species (Lycopersicon esculentum). Agriculture Water Management, 2005, 78:128-135.
- [33] W. Yu-feng, Effect of NaCl stress on seed germination of watermelon. Journal of Anhui Agriculture Science, 2006, 34(24):6497-6499.

Citation: Cenk PASA. Different Nacl Concentrations Impact on Seed Germination of Basil (Ocimum Basilicum L.) and Fennel (Foeniculum VulgareL.) Cultivars. International Journal of Advanced Research in Botany. 2025; 9(1):1-4. DOI: https://doi.org/10.20431/2455-4316.0901001.

Copyright: © 2025 Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.